


Chapter 5

The Active Sun



5 The Active Sun

Other than the solar cycle itself, all of the phenomena that affect HF communications occur in the Sun's atmosphere. This is particularly true during solar maximum. These phenomena include

- Sunspots,
- Faculae,
- Plages,
- Prominences,
- Filaments,
- Solar Flares,
- Coronal Mass Ejections (CMEs),
- Coronal Holes, and
- Radio Noise.

5.1 Sunspots and Sunspot Groups

Sunspots, as we know, are black irregularly shaped blemishes on the Sun's surface. A number of sunspots are shown in Figure 1. Sunspots appear, increase in number, and then gradually disappear over the 11 year solar cycle. A solar cycle is arbitrarily defined to begin at solar minimum when few if any sunspots are present. Increasing numbers of sunspots appear at a fairly rapid rate as the cycle progresses. The solar cycle reaches a maximum in roughly 3 to 6 years when a relatively large number of sunspots are visible. The sunspots then slowly disappear over the next 5 to 8 years reaching a minimum again at the end of the cycle.



Figure 1 Sunspot groups (credit: NASA Goddard Space Flight Center)

A sunspot develops at a point where the Sun's magnetic field erupts through the photosphere (the Sun's visible surface) as shown in Figure 2. A second sunspot occurs where the magnetic field plunges back into the photosphere. Notice that sunspots have magnetic polarities created by the magnetic field's direction of flow. The field flows out of a magnetically north (N) sunspot and into a south (S) magnetic spot. North and south are often represented as "+" and "-" respectively.




Figure 2 Formation of sunspots (credit: Sky Maps with Pierre Auger Data)

The peak intensity of the field emanating from a sunspot is in the neighborhood of 3,000 to 4,000 gauss. This field is so intense that it suppresses the upward flow of hot plasma below the sunspot location, causing the sunspot site to be over 2,000 degrees cooler than the surrounding photosphere. The sunspot's lower temperature is why it appears black in color.

The center of a sunspot is the umbra while the outer lighter colored region is the penumbra shown in Figure 3. The umbra is the lowest temperature (blackest) part of a sunspot. The temperature of the umbra usually ranges from about 3,800 to 4,100 degrees kelvin compared to 6,500 K at the base of the photosphere. The penumbra around the outer edge of the sunspot is warmer and lighter in color. It is characterized by large numbers of elongated dark and bright filaments that extend outward from the umbra. Out flows of material occur in the penumbra, beginning at the umbra, reaching a maximum speed of 2 – 6 km/s, and then dissipating outside the penumbra.

Sunspots can be huge. Large sunspots (including both the umbra and penumbra) can be 50,000 km or more in diameter. More typically, the umbra is around 10,000 km in diameter, about the same size as the Earth. The Earth's diameter is 12,756 km. Sunspots appear, last several days, and then disappear. Some sunspots may last for several weeks.

Sunspots are part of the Sun's photosphere and thus rotate with the Sun. In Figure 4 sunspots move from left to right as the Sun rotates. Sunspots that disappear around the Sun's right limb may reappear about 13 days later on the left side of the Sun.




Figure 3 A closer look at sunspots (credit: spaceweatherlive.com)




Figure 4 Sunspots on the face of the Sun (credit: NOAA Space Weather Prediction Center)

While isolated individual sunspots do form, most sunspots occur in pairs, or more commonly in sunspot groups aligned with the Sun's direction of rotation. A typical sunspot group is shown in Figure 5. The group rotates with the Sun so the large sunspots on the right are the leading spots while those on the left are the trailing or following sunspots.




Figure 5 Sunspot Group (source: SpaceWeatherLive.com)

Sunspot groups tend to be complex varying widely in size and configuration. Considerable differences not only occur between groups but within the same group as it evolves.

There are important distinctions between leading and following sunspots. Leading sunspots are nearly always larger than following spots. Leading sunspots also tend to be more stable and have a longer life. Following sunspots seldom achieve long-term stability and often break up within a few days after appearing. Following sunspots also tend to be more irregular in shape often having only partial penumbras. 95% of sunspot groups that last long enough to make a second pass across the solar disk do so only with a single leading sunspot.

The leading and following sunspots have opposite magnetic polarities. If the leading sunspot has a north magnetic polarity, then the following sunspots must have south polarities. In addition, the leading sunspot always has the same magnetic polarity as the hemisphere in which it is located. The leading sunspot will have a north magnetic polarity in the Sun's magnetically northern hemisphere. Consequently, the leading sunspots in the northern and southern hemisphere have opposite polarities. The same is true of the following sunspots. Because of this, sunspot groups on opposite sides of the solar equator cancel each other out resulting in no sunspots at the equator itself.

Sunspot groups are inclined with the leading sunspot being closer to the equator than the following spots. You can see this inclination in Figure 5. In this figure the leading spot is closer to the bottom of the picture, in the direction of the equator, than the other sunspots. The inclination is illustrated in more detail in Figure 6. The amount of inclination varies with latitude. Near the equator the inclination is usually only a few degrees. At higher latitudes the inclination may be as much as 15° .




Figure 6 Inclination of a Sunspot Group (credit: author)

This inclination, coupled with the Sun's differential rotation, cause sunspots within a group to move with respect to one another. This movement tends to twist prominences that are rooted in sunspot groups. If the twisting is great enough it becomes one of the factors leading to a solar flare.

5.2 Faculae

Faculae are bright hot depressions in the photosphere at locations where strong vertical magnetic fields emerge from the Sun's surface. However, a faculae magnetic field is not nearly as strong as that found in the umbra of a sunspot. Faculae are hotter than the surrounding photosphere which is why they appear bright. Within a faculae heat radiates from the walls of the faculae depression as well as from the bottom of the depression accounting in part for the faculae being hotter and brighter than the surrounding photosphere.

About 90% of faculae are associated with sunspots. Faculae typically appear several days before the appearance of sunspots and survive considerably longer. In terms of HF radio propagation, faculae are important because they are associated with hot Extreme Ultra Violet (EUV) emitting plages located in the chromosphere above the faculae.

5.3 Plages

Plages are hot bright irregularly shaped areas easily visible in the image of the chromosphere shown in Figure 7. Plages are formed in the chromosphere by intense magnetic fields radiating out from faculae in the underlying photosphere. Like the faculae below them, plages occur in active sunspot regions of the Sun and usually form several days prior to sunspots in the area. Plages typically last long periods of time, longer than their associated sunspots, and emit copious amounts of EUV radiation responsible for ionizing Earth's upper atmosphere.




Figure 7 Sun's chromosphere shown in H alpha (H_{α}) light (credit: universitytoday.com)

While there is a strong correlation between sunspots and long distance radio communications, it turns out that sunspots have little to do with HF propagation. Sunspots are far too low in temperature to generate the EUV radiation needed to ionize Earth's upper atmosphere. The required EUV radiation is primarily produced by plages. The problem is that plages can not be seen in the

intense light emanating from the photosphere because the photosphere is too bright. But sunspots can easily be seen as readily apparent in Figure 1. Thus sunspots become markers for plages. A large number of sunspots means a large number of plages, high EUV levels, strong ionization of the Earth's ionosphere, and good HF radio communications.

The EUV radiation responsible for ionizing Earth's upper atmosphere accounts for only 0.001% of the Sun's total energy output as illustrated in Figure 8. While the Sun's energy output remains incredibly stable over thousands of years, the Sun's EUV energy output is NOT stable at all! EUV energy varies considerably over the 11 year solar cycle. Consequently the level of ionization in Earth's upper atmosphere also varies over the same time period. EUV energy output is greatest during solar maximum when there are large numbers of sunspots, and consequently large numbers of EUV emitting plages. EUV output is lowest during solar minimum when there are few if any sunspots and only a few randomly scattered plages.




Figure 8 Sun's energy output (credit: author)

5.4 Prominences and Filaments

Prominences are large bright loops and curtains of relatively cool plasma suspended high above the photosphere by strong arching magnetic fields. A prominence occurring on the edge of the Sun is shown in Figure 9.




Figure 9 Active Region Eruptive Solar Prominences (credit: NASA)

A prominence occurs when the Sun's stressed magnetic field erupts through the photosphere carrying with it electrically charged plasma. The charged plasma, often traveling at speeds of 40 km/sec, follows the contours of the magnetic field giving a prominence its bright arching shape. The strength of the magnetic field within a prominence typically ranges from 10 to 800 gauss, compared to the Sun's quiet bipolar field of around 1 gauss and intensities of over 3,000 gauss in magnetic field lines that become twisted and knotted. A prominence extends deep into the corona often 50,000 km or more above the photosphere and usually lasts for several hours. In some cases a prominence can remain in place for 2 - 3 solar rotations. The temperature within a prominence is on the order of 7,000 degrees kelvin, much cooler and more dense than the surrounding corona. The stability of a prominence is due to the equilibrium between its opposing magnetic and gravitational forces. Disruption of this equilibrium causes prominences to collapse, sometimes catastrophically producing solar flares and coronal mass ejections.

Filaments and prominences are the same thing seen from different perspectives. A prominence appears as a high bright arching formation when viewed on the edge of the Sun against the black background of interstellar space. When viewed on the face of the Sun, it appears as a dark erratic scar or filament, as illustrated in Figure 10. A filament is dark in appearance because it is relatively cool compared to the chromosphere below it.




Figure 10 Filaments on the solar disk in H – alpha light. (credit: Big Bear Solar Observatory)

Prominences appear in various shapes and sizes. Prominences occurring in active regions of the Sun include eruptive prominences, like the one shown in Figure 9, spray prominences shown in Figure 11 that often precede a solar flare, and loop prominences shown in Figure 12 that typically form after a flare. In active regions prominence material is often observed condensing and falling back toward the photosphere forming falling prominences like the one shown in Figure 13. Falling prominences are sometimes referred to as coronal rain.

Quiescent prominences are typically found far from active regions of the Sun. The hedgerow prominence shown in Figure 14 is a common quiescent prominence.




Figure 11 Pre-flare Spray Prominence (credit: cloudy nights.com)




Figure 12 Post Flare Loop Prominence (source: Astronomy Magazine)




Figure 13 Falling “coronal rain” Prominence (credit: space.com)



Figure 14 Quiescent hedgerow prominence (Source: Sky & Telescope)

5.5 Solar Flares

In active regions of the Sun arching magnetic fields rooted in the photosphere extend high into the corona creating coronal loops (prominences). Figure 15 shows a large coronal loop with Earth superimposed for reference. Hot plasma flowing along the magnetic field lines give the coronal loop its bright arching appearance.

A solar flare is a massive, sudden, explosive releases of energy stored in the magnetic field of a coronal loop. A flare occurs when the magnetic field lines of a coronal loop become severely stretched out and twisted concentrating enormous magnetic energy in a relatively small confined region of the loop. Continued stretching and twisting squeeze magnetic field lines closer and closer together. As this happens, energy in the confined region builds until it becomes so intense that outbound (+) and returning (-) magnetic field lines “short circuit” releasing a massive amount of

energy in the form of a flare, a flash of light. In the process the outbound and returning magnetic field lines connect at the rupture forming a smaller lower energy magnetic loop, still anchored in the photosphere, that is only a tiny fraction of its original size. The top part of the initial coronal loop is blown off into interstellar space.




Figure 15 Coronal loop (Image credit: NASA's Cosmos)

A flare last from a few minutes to several hours heating the corona plasma in the vicinity of the explosion to well over ten million degrees. Extensive heating also occurs in the underlying chromosphere and in some cases in the photosphere. Large solar flares frequently occur during solar maximum and are the most powerful eruptions in the solar system.

A large solar flare occurring on Oct 2, 2014, captured by NASA's Solar Dynamics Observatory, is shown in Figure 16. This image of the chromosphere was taken in a hydrogen-alpha (H_α) light. The flare is the large bright flash of light seen just above the chromosphere near the edge of the Sun. Below the flare a burst of solar material is erupting out into space. Flares are rarely visible in normal light because the photosphere is too bright.

Solar flares emit strong bursts of electromagnetic radiation, including gamma rays, X-rays, extreme ultraviolet radiation, visible light, and radio waves. In addition, huge quantities of high energy particles travel outward at high speeds from solar flares. Solar flares may also eject large clouds of coronal and chromospheric material, known as a coronal mass ejection (CME), into space and generate powerful magnetic shock waves. The shock waves propagate at high speed ahead of the CME material deep into interstellar space seriously disrupting magnetic fields well beyond Earth's orbit.

The X-ray and EUV radiation from a solar flare reach the Earth in about 8 minutes. High energy particles can arrive in as little as 30 minutes. Shock waves and coronal mass ejections arrive in 1 to 4 days.




Figure 16 Solar flare (credit: NASA/SDO)

A coronal loop begins in Figure 17 as a strong arching magnetic field between a leading and following sunspot. As discussed earlier, sunspot groups are inclined with the leading spot closer to the equator than the following spots. Differential rotation of the Sun's surface coupled with convection zone turbulence cause spots in a sunspot group to move relative to one another, twisting the coronal loop illustrated in Figure 18 as it stretches out high into the corona. Continued stretching and twisting cause outgoing and returning magnetic field lines to come in close proximity at a region of the loop known as the x – point, so named because of its shape. This is shown in Figure 19. Tremendous magnetic forces build up at the x – point eventually causing the coronal loop to rupture at that point releasing enormous energy in a massive solar flare explosion. The outgoing and returning magnetic fields reconnect at the x – point (#1 in Figure 20) forming a smaller lower energy loop below the rupture – a process called magnetic reconnection. This smaller loop is rooted in the photosphere. A second loop forms above the rupture extending outward away from the Sun as illustrated in Figure 20.



Figure 17 Emerging coronal loop (credit: UC Berkeley)




Figure 18 Coronal looping stretching out high into the corona (credit: UC Berkeley)




Figure 19 X-point forming (credit: UC Berkeley)

Below the rupture hot plasma consisting of energetic electrons plus hydrogen and helium nuclei stream down the reconnected field lines (#3 in Figure 20). These energetic particles are traveling at nearly the speed of light. Intense heat is generated as they crash into the upper chromosphere. In some cases the intense heat and relativistic speed of the collisions cause sporadic nuclear reactions to occur within the chromosphere releasing gamma rays and hard X-rays in addition to EUV radiation.

Above the rupture a hot plasma, also consisting of energetic electrons plus hydrogen and helium nuclei, stream outward along the upper field lines into interstellar space (#2 in Figure 20). The kinetic energy associated with the rapidly expanding plasma creates shock waves (#4 in Figure 20) that propagate out from the Sun far past Earth's orbit.




Figure 20 Structure of a solar flare (credit: ResearchGate)

5.6 Coronal Mass Ejections (CMEs)

A coronal mass ejection (CME) is huge eruption of coronal plasma, with its associated frozen-in magnetic field, that moves outward from the Sun into interplanetary space. CMEs vary widely in size, shape, and speed. Some look like loops, other like bubbles, and some are irregular in shape. Figure 21 is an example of a CME.

CMEs are often caused by the catastrophic collapse of coronal loops as magnetic fields re-align and reconnect into lower energy states. CMEs may also be triggered by solar flares.

A CME can eject billions of tons of coronal material outward from the Sun at speeds typically ranging from 200 to 500 km/s. Some energetic CMEs can reach speeds of 3,000 km/s or more. A shock wave is created when the CME travels faster than the background solar wind speed. The shock wave often accelerates charged particles ahead of it which can create intense radiation storms as the CME impacts the Earth. During solar maximum several CMEs of various sizes and shapes occur per day. At solar minimum one CME is typically observed every 5 days or so.




Figure 21 Coronal Mass Ejection (credit: www.astronet.ru)

Most CMEs are ejected outward from the Sun into the solar system away from the Earth. However, a CME launched in the direction of Earth can arrive in as little as 15-18 hours. Slower CMEs may take several days to arrive. A CME expands as it travels away from the Sun. A large CME can encompass nearly a quarter of the distance between the Sun and Earth by the time it reaches Earth.

Forecasters at the NOAA Space Weather Prediction Center (SWPC) monitor CMEs using imagery from orbital satellites to determine the likelihood of a CME impacting Earth. These satellites utilize an instrument known as a coronagraph to block out the Sun's photosphere so that the corona can be observed. A coronagraph in effect creates an artificial solar eclipse as shown in Figure 21. The NASA Solar and Heliospheric Observatory (SOHO) carries a coronagraph – known as the Large Angle and Spectrometric Coronagraph (LASCO). This instrument has two ranges for optical imaging of the Sun's corona: C2 (covers distance range of 1.5 to 6 solar radii) and C3 (range of 3 to 32 solar radii). The LASCO instrument is currently the primary means used by forecasters to analyze and categorize CMEs; however another coronagraph is on the NASA STEREO-A spacecraft as an additional source.

The SOHO and Deep Space Climate Observatory (DSCOVR) spacecrafts both orbit the Sun at the L1 point approximately 1 million miles away from Earth. The L1, or Lagrangian Point 1, is the neutral gravity point between the Earth and the Sun and provides an uninterrupted view of the Sun.

A sudden increases in particle density, interplanetary magnetic field (IMF) strength, and solar wind speed at the DSCOVR spacecraft indicates an approaching CME shock wave and associated plasma cloud with its frozen-in magnetic field. This early detection provides a 15 to 60 minute warning prior to the CME impacting the Earth and potentially triggering sudden geomagnetic storms.

5.7 Coronal Holes

Coronal holes are dark areas seen in the solar corona when viewed in EUV and soft X-ray light. They appear dark, like those shown in Figure 22, because they are cooler and less dense than the surrounding corona plasma.

Coronal holes can develop at any time and any location on the Sun. However, they occur most often at the solar north and south poles. Some of these grow and expand into the lower latitudes. Coronal holes can also develop in other regions of the Sun independent of the polar holes. In some cases independent holes split off from polar holes. Coronal holes occur most often and last longer during solar minimums. In some cases coronal holes can last for several solar rotations

Coronal holes are regions of open, unipolar magnetic fields that allow streams of relatively fast solar winds, known as high speed streams, to easily escape into space. In fact, long-lasting coronal holes are a major source of high speed solar wind streams that can buffet Earth for many days. Space weather forecasters analyze coronal holes closely because of their potential to disrupt Earth's magnetic field creating geomagnetic storms.




Figure 22 Coronal Holes (credit: NOAA www.swpc.noaa.gov/phenomena)

5.8 Solar Radio Noise

The Sun produces various forms of radio noise including:

1. Continuous background noise that is always present, and
2. Radio bursts that occur following flares and other disruptive events on the Sun.

5.8.1 Background Solar Radio Noise

The Sun produces continuous background radio noise generated by random collisions of electrons with heavier particles high in the chromosphere and lower corona. This radio noise is easily

measured at a frequency of 2,800 MHz (10.7 cm wavelength) and is known as the Solar Flux Index (SFI). It has been consistently measured on a daily basis in Canada since 1947, first in Ottawa, Ontario and more recently at the Penticton Radio Observatory in British Columbia. It is one of the longest running records of solar activity. Because this radio noise originates in the Sun's chromosphere and corona, it tracks other important emissions that form in the same regions of the solar atmosphere. The Solar Flux Index correlates well with measured levels of solar X-ray and Extreme Ultra Violet (EUV) radiation. It also correlates remarkably well with sunspot numbers. In addition, the Solar Flux Index is an excellent indicator of ionospheric conditions, including HF radio propagation through Earth's upper atmosphere. SFI numbers vary from a low of about 65 during solar minimum to around 225 at the peak of a solar cycle.

5.8.2 Radio Bursts

Figure 23 illustrates four types of radio bursts that commonly occur:

- Type I: Noise storm
- Type II: Slow drift burst
- Type III: Fast drift burst
- Type IV: Prolonged continuum




Figure 23 Solar radio bursts (Credit: ResearchGate)

The bursts are classified based on their spectrum, duration, and cause.

5.8.2.1 Type I Noise Storm

A Type I noise storm is a slowly varying increase in the Sun's background radio noise. The increase may be a million times greater than normal background conditions. Type I storms are strongly circularly polarized with a relatively narrow bandwidth typically ranging in frequency from 100 to 400 MHz. A storm can last from several hours to several days and may drift higher or lower in frequency as it evolves. Type I storms originate in the corona above active flare producing regions of the Sun.

5.8.2.2 Type II Slow Drift Radio Bursts

A Type II radio burst is an intense release of RF energy at frequencies below 200 MHz. The burst occurs in two discrete frequency bands consisting of fundamental and second harmonic frequencies. The two frequency bands drift lower in frequency at a rate of around 1 MHz per second over a period of 5 to 20 minutes. As shown in Figure 23, Type II bursts are generated by plasma disturbances (shock waves) that typically occur around the time of the soft X-ray peak in a solar flare. These disturbance expand outward from active regions of the Sun at 500 to 5,000 km/sec. The slow downward shift in frequency is the result of the disturbances traveling through regions of the corona having progressively lower electron densities.

5.8.2.3 Type III Fast Drift Radio Bursts

Type III radio bursts occur during the initial phase of solar flares and are the most common type of burst. They generate RF energy at frequencies from 50 KHz to around 700 MHz. They drift lower in frequency also, but much faster than Type II. The drifting rate for a Type III burst is around 10 - 20 MHz per second. Type III bursts are much shorter than Type II lasting only a few seconds to a few minutes. They are generated by plasma disturbances caused by energetic electrons accelerating outward from active regions of the Sun at speeds of 100,000 km/sec, as shown in Figure 23. The rapid downward shift in frequency is the result of these electrons traveling at high speeds through regions of the Corona with progressively lower electron densities.

A Type III noise storm is the decaying phase of a Type III burst that occurred earlier. A Type III storm is a low frequency long duration event as shown in Figure 23.

5.8.2.4 Type IV Prolonged Radio Bursts

Type IV radio bursts are quite different from Type II and III. They are generally considered the radio signature of coronal mass ejections (CME's) produced by flares and erupting coronal loops. They produce intense RF energy at frequencies from 100 KHz well into the microwave range. However, the most studied Type IV bursts occur in the range from 10 to 100 MHz. Type IV bursts typically last from 10 minutes to an hour or more and are believed to be generated by gyro-

synchrotron radiation from energetic electrons magnetically trapped in plasma clouds. These clouds typically accelerate outward from solar flares at speeds from a few hundred to about 1,500 km/sec.

5.9 Detection of Solar Flares

A solar flare is detected by a rapid increase in solar X-ray energy from the Sun. The increase occurs in minutes followed by a gradual decline over tens of minutes to several hours. Figure 24 is a large flare captured by a Soviet Union spacecraft on September 7, 2005. Figure 25 is the X-ray flux measured by the GOES 12 spacecraft of the same flare.




Figure 24 Image of the solar flare occurring on September 7, 2005 taken by the Soviet Union Spirit spacecraft.

The magnitude of solar flares are classified by their peak X-ray flux in watts per square meter on a logarithmic scale. The scale consists of 5 Classes.

- Class - A: typical background X-ray flux levels.
- Class - B: elevated background X-ray flux levels.
- Class - C: small minor flare with few noticeable affects on Earth.
- Class - M: medium size flare that can cause brief HF radio blackouts. The blackouts may be followed by minor radiation storms.
- Class - X: major flare event that can cause large scale HF radio blackouts and long lasting radiation storms




Figure 25 Solar X-ray flux measured by the GOES 10 and 12 spacecraft on September 7, 2005

Each class has a peak flux ten times greater than the preceding class. That is a Class M flare is ten times more powerful than a Class C. The classification of flares is thus similar to the measurement of earthquakes on the Richter scale. A magnitude 6 earthquake is 10 times stronger than a magnitude 5 quake. Each class of X-ray flux is in turn divided into nine logarithmic subclasses, for example B1 through B9. The next level above B9 is C1.

The X-ray flux resulting from the flare in Figure 25 went from a normal background level of A3 (as measured by the GOES 12 spacecraft in the 0.5 – 4.0A spectrum) to X5 in about 10 minutes. The graph in Figure 25 was produced 3 hours later. By then the X-ray flux had dropped to around C2. However, it was several more hours before the X-ray flux returned to Class B levels.

The GOES (Geostationary Operational Environmental Satellite) series of spacecraft are operated by NOAA. A drawing of the GOES 16 spacecraft is shown in Figure 26. The GOES satellites continuously measure X-ray and EUV radiation from the Sun. In addition the GOES satellites measure high energy proton (hydrogen nuclei) and alpha particle (helium nuclei) densities at the satellite's location in geosynchronous orbit around the Earth. The GOES satellites also measure

infrared radiant energy and visible reflected solar energy from the Earth's surface and atmosphere including vertical atmospheric temperature and moisture profiles, surface and cloud top temperature, and ozone distribution.




Figure 26 GOES 16 Spacecraft (Source: NOAA)

The Earth's ionosphere reacts badly to large flares. Figure 27 shows the attenuation of radio frequencies resulting from the September 07, 2005 flare. The attenuation is caused by the flare's X-ray energy heavily ionizing the ionosphere's D – Layer. The attenuation is greatest at low frequencies and is progressively less at higher frequencies. This is clearly indicated by the red bar graph at the right side of Figure 27. Radio black out conditions begin at low frequencies and rapidly extend to higher frequencies. For this particular flare all frequencies up through 20 MHz were blacked out. The position of the Sun is shown in this figure by the black diamond. Notice that regions of the Earth directly below the Sun (at noon local time) are impacted the most. But for this particular flare, all of North and South America experienced radio black outs on all frequencies below 20 MHz. This figure also shows that the expected recovery time from this flare was 1 hour 17 minutes. The estimate was drastically in error. The flare event last for several hours, eventually blacking out most of East Asia.

This solar flare is particularly interesting in that the author and two colleagues, quite by accident, noticed and began tracking the flare the moment that it occurred. We had been participating in the weekly California Emergency Services Net on 40 meters. Prior to the net we recorded the solar X-ray flux and critical frequency as was our procedure. We also recorded the signal strengths of WWV (Boulder Colorado) and WWVH (Kauai, Hawaii) on 5, 10, 15, and 20 MHz. The signals on 5, 10, and 15 MHz were strong. 20 MHz was there but weak. The net went well. We heard most stations throughout California and they heard us. The net closed as normal. Immediately following the net we checked WWV - WWVH signal strengths. 5 MHz was completely gone. There were no signals on 5 MHz. Signals on 10 MHz were very erratic rapidly fading in and out. Reception on 15 MHz was still good with WWV and WWVH both coming in strong. 20 MHz was still there but weak as before. We stepped back down in frequency. 15 MHz, 10 MHz, and 5 MHz were all gone. No signals on these frequencies. We went back up to 20 MHz. It too was gone. We concluded that something was badly wrong and guessed that it was probably a flare. We check the NOAA X-ray flux levels and watched the level rise rapidly to Class X while radio black out expanded in frequency. It was an exciting event to experience.




Figure 27 Attenuation of radio signals resulting from the September 07, 2005 flare

References

Foukal, Peter; “Solar Astrophysics third edition”; Wiley-VCH Publishing Company, 2013

Carroll, Bradley W. and Ostlie, Dale A.; “An Introduction to Modern Astrophysics”; Addison-Wesley Publishing Company Inc., 1996

Golub, Leon and Pasachoff, Jay M.; “Nearest Star The Surprising Science of Our Sun second edition”; Cambridge University Press, 2014

Moldwin, Mark; “An Introduction to Space Weather”; Cambridge University Press, 2008

White, Stephen M.; “Solar Radio Bursts and Space Weather”; Dept. of Astronomy, University of Maryland, College Park, MD 20742 USA

Friedman, Herbert; “The Astronomer’s Universe Stars, Galaxies, and Cosmos”; W. W. Norton & Company, 1990

Davies, Kenneth; “Ionospheric Radio”; Peter Peregrinus Ltd., 1990

Levis, Curt A. ; Johnson, Joel T.; and Teixeira, Fernando L.; “Radiowave Propagation Physics and Applications”; John Wiley & Sons, Inc., 2010

McNamara, Leo F.; “The Ionosphere: Communications, Surveillance, and Direction Finding”; Krieger Publishing Company, 1991

DeSoto, Clinton B., “200 Meters & Down”; The American Radio Relay League, Inc., 1936

Yeang, Chen-Pang; “Probing The Sky With Radio Waves”; The University of Chicago Press, 2013

Piccioni, Robert L.; Everyone’s guide to Atoms Einstein and the Universe second edition”; Real Science Publishing, 2010

Hogan, Craig J.; “The Little Book of the Big Bang A Cosmic Primer”; Springer-Verlag New York, Inc., 1998

Zirin, Harold and Lang, Kenneth; “Sun - Astronomy”; Encyclopedia Britannica

Kawaler, Steven D., and Dahlstrom, Michael; “White Dwarf Stars”; American Scientist, 2000 [pdf]

“White Dwarf Stars” National Aeronautics and Space Administration, Goddard Space Flight Center, 2010

“Solar experts predict the Sun’s activity in Solar Cycle 25 to be below average, similar to Solar Cycle 24”; 2019 NOAA Space Weather Workshop in Boulder, Colo.

Dobler, Sacha; “The next Grand Solar Minimum has (very likely) begun: NASA predicts lowest solar cycle in 200 years”; 14 June 2019

Kuznetsov, V. D.; “From the geophysical to heliophysical year: The results of the CORONAS-F project”; Russian Journal of Earth Sciences, Vol. 9, ES3004, doi: 10.2205/2007ES000276, 2007

Hathaway, David H., Wilson, Robert M., Reichmann, Edwin J.; “Group Sunspot Numbers: Sunspot Cycle Characteristics”; NASA/Marshall Space Flight Center, Huntsville, AL., 2002

Minzner, R. A.; “The 1976 Standard Atmosphere Above 86 km Altitude” NASA Goddard Space Flight Center, 1976

Masi, Marco; “Quantum Physics: an overview of a weird world”; Marco Masi 2018

Zettili, Nouredine; “Quantum Mechanics Concepts and Applications second edition”; John Wiley & Sons, 2009

Ahrens, C. Donald; “Essentials of Meteorology”; Wadsworth Publishing Company, 1998

“Aviation Weather”; Department of Transportation FAA and Department of Commerce NOAA, 1975

Williams, Matt; “What is heat conduction?”; Universe Today, 12/9/2014

Halliday, David and Resnick, Robert; “Physics For Students of Science and Engineering Part II second edition”; John Wiley & Sons, Inc., 1962

Zumdahl, Steven S. “Chemistry second edition”; D. C. Heath and Company, 1989

Devoldere, John; “Low-Band DXing” fourth edition; ARRL, 2005

“The ARRL Antenna Book For Radio Communications”; ARRL

Whitfield, Philip; “From So Simple A Beginning”; Marshall Editions Developments Limited, 1993

Roberts, Alice; “Evolution The Human Story”; DK, 2011

Potts, Richard and Sloan, Christopher; “What Does It Mean To Be Human?”; National Geographic,

Dr. David Hathaway, Dr. Lisa Upton "Solar Cycle Science Discover the Sun!"
<http://solarecyclescience.com/>

March 2020

Kenneth N. Larson KJ6RZ

Solar Physics Group at NASA's Marshall Space Flight Center
<https://solarscience.msfc.nasa.gov>